

YOU HAVE BEEN PRESCRIBED


Pr **VYZULTA**™

Latanoprostene
bunod ophthalmic
solution, 0.024% w/v

to reduce the high pressure
in the eye for patients with
open-angle glaucoma or
ocular hypertension



YOU ARE NOT ALONE!

In fact, more than
250,000 Canadians have
open-angle glaucoma.

Speak to your doctor
for more information about
your condition.

TAKING VYZULTA™



USUAL DOSE

- The recommended dose of VYZULTA™ is one drop into the affected eye(s) **once daily** in the **evening**.
- Apply **one** drop of VYZULTA™ into each affected eye.
- Use VYZULTA™ exactly as your doctor has directed. Do not apply VYZULTA™ more than once a day.
- If your doctor has prescribed other eye drops to be taken along with VYZULTA™, apply each medicine at least 5 minutes apart.



Set a reminder to take your
drops at the same time each day.



MISSED DOSE AND OVERDOSE

- If you forget to use VYZULTA™, continue your treatment with the next dose the following day.
- If you think you have taken too much VYZULTA™, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.



REMEMBER

- Be careful not to touch your eye, eyelids, surrounding areas, fingers, or any other surface with the dropper tip of the bottle. If you do, you may contaminate the medicine, and this can cause you to have an eye infection.



- Avoid touching the tip of the bottle.
- If you wear contact lenses, remove them before applying VYZULTA™.
- You may reinsert your lenses 15 minutes after you have applied VYZULTA™.



It is **very important** that you take VYZULTA™ **exactly as directed** by your healthcare professional. If you have any questions, talk to your healthcare professional.



For more information about VYZULTA™, please talk to your healthcare professional and consult the patient medication information leaflet that came with your medication.



STORAGE

- Before VYZULTA™ is first opened, keep it in the fridge (between 2°C and 8°C).
 - Once the VYZULTA™ bottle has been opened, it can be stored at room temperature (up to 25°C).
 - VYZULTA™ should not be used after the expiry date on the bottle.
 - **Protect from light and freezing.**
 - Keep out of reach and sight of children.
- Discard the bottle and/or unused content after 8 weeks.



WARNINGS AND PRECAUTIONS

Do not use VYZULTA™ if you:

- are allergic to latanoprostene bunod (the medicinal ingredient in VYZULTA™)
- are allergic to any of the other ingredients in VYZULTA™ or to a component of the container

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take VYZULTA™.

Talk about any health conditions or problems you may have, including if you:

- have or have had eye inflammation including the conditions iritis or uveitis
- have or have had a torn posterior lens capsule
- have or have had eye swelling including a condition called macular edema
- have or have had a viral infection of the eye called herpes keratitis
- have severe asthma or asthma that is not controlled
- are using any other eye drops
- notice any changes in your eyelashes or colour of your eyes
- wear contact lenses
- are pregnant or planning to become pregnant
- are breastfeeding or planning to breastfeed

Other warnings you should know about:

- Driving and using machines
 - Wait until you can see clearly before driving or operating machines after applying VYZULTA™.
- Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

POSSIBLE SIDE EFFECTS

Common side effects

- eye irritation
- eye redness
- eye pain
- pain in your eye at the application site
- runny or stuffed nose, sore throat

Rare side effects

- altered sense of taste
- feeling of grittiness or having something in the eye

Other side effects

- VYZULTA™ may change colour of your eye or eyelid. The eye colour change may not be noticeable for many months to years. The eye colour and eyelid changes may be permanent. Tell your doctor if your eye colour has changed. They may examine your eyes more often.
- VYZULTA™ may also cause your eyelashes to appear thicker and longer than they usually do and increase in number. Eyelash changes are reversible after treatment with VYZULTA™ is stopped.

SERIOUS SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Talk to your healthcare professional in all cases if you develop the following rare side effects:

- Bacterial keratitis (infection of the eye): eye pain or redness, decreased vision, discharge from the eye, sensitivity of the eye to light, swelling of the eye or eyelid, watery eyes or production of tears.
- Macular edema (swelling and buildup of fluid in the center of the retina): blurry vision, blurry or wavy vision near or in the center of your field of vision, colours may appear washed out or faded.

These are not all the possible side effects that you may feel when taking VYZULTA™. If you experience any side effects not listed here, contact your healthcare professional.

Talk to your healthcare professional if you want more information about VYZULTA™. Find the full Product Monograph that is prepared for healthcare professionals and includes the Patient Medication Information leaflet by scanning the QR code below. The Product Monograph is also available by calling **1-888-459-5000**.

